

Syphilis cases are up in B.C.

You may have it and not know it. You can spread it, even if you don't have symptoms.

Syphilis can be treated with antibiotics, treatment is free. If left untreated, syphilis can be serious.

Syphilis testing is an important part of health care. If you're between the ages of 18-55, syphilis testing will be added to your bloodwork today.

Speak with our staff if you have any questions.



Speak with our staff if you have any questions.

Scan here to learn more: vch.ca/SyphilisTesting







Syphilis rates are rising in BC, especially among under-housed populations and patients with substance use disorders aged 18 to 59.

Test high-risk patients every 3-6 months and patients who are symptomatic, regardless of risk, using serology.

PCR testing does not replace serology but should be collected if individuals present with symptoms consistent with primary or secondary syphilis (use an Aptima/viral swab for TP PCR).



Speak with our staff if you have any questions.

Scan here to learn more: vch.ca/SyphilisTesting







Syphilis testing is an important part of healthcare.

Syphilis rates are rising in BC, especially among underhoused populations and patients with substance use disorders aged 18 to 59.

Consider testing high-risk patients every 3-6 months and patients who are symptomatic, regardless of risk, using serology.

PCR testing does not replace serology but should be collected if individuals present with symptoms consistent with primary or secondary syphilis (use an Aptima/viral swab for TP PCR)



Speak with our staff if you have any questions.

Scan here to learn more: vch.ca/SyphilisTesting



